

LOWER BOUND FOR THE GEOMETRIC TYPE FROM A GENERALIZED ESTIMATE IN THE $\bar{\partial}$ -NEUMANN PROBLEM - A NEW APPROACH BY PEAK FUNCTIONS

TRAN VU KHANH

ABSTRACT. We give a simple proof of the fact that an “ f -estimate” for the $\bar{\partial}$ -Neumann problem implies a lower bound on the geometric type of the boundary along any complex one dimensional variety. The proof uses the existence of peak functions which is in turn a consequence of the f -estimate.

1. INTRODUCTION

In a series of seminal papers in the Annal of Mathematics [Cat83, Cat87], Catlin proved the equivalence of the finite type of a boundary (cf. [D’A82]) with the existence of a subelliptic estimate for the $\bar{\partial}$ -Neumann problem by triangulating through the t^ϵ -Property (see below)

- (i) finite type $m \Rightarrow t^\epsilon$ -Property with $\epsilon = m^{-n^2 m^{n^2}}$;
- (ii) t^ϵ -Property $\Rightarrow \epsilon$ -subelliptic estimate;
- (iii) ϵ -subelliptic estimate $\Rightarrow$ finite type m for $m \leq \frac{1}{\epsilon}$.

Here, the t^ϵ -Property of a boundary $b\Omega$ is a case of a more general “ f -Property” defined as follows. For a smooth monontonic increasing function $f : [1, +\infty) \rightarrow [1, +\infty)$ with $f(t) \leq t^{\frac{1}{2}}$, the f -Property at z_o consists in the existence of a neighborhood U of z_o , of constant C_1, C_2 and of a family of functions $\{\phi_\delta\}$ such that

- 1) ϕ_δ are plurisubharmonic and C^2 on U and $-1 \leq \phi_\delta \leq 0$;
- 2) $\partial\bar{\partial}\phi_\delta \geq C_1 f(\delta^{-1})^2 Id$ and $|D\phi_\delta| \leq C_2 \delta^{-1}$ for any $z \in U \cap \{z \in \Omega : -\delta < r(z) < 0\}$, where r is a defining function of Ω .

Also, the finite type m is a case of a general “ F -type” along a one-dimensional complex analytic variety Z consisting in the estimate

$$|r(z)| \leq F(|z - z_o|) \quad z \in Z, z \rightarrow z_o. \quad (1.1)$$

The results in the step (ii) and (iii) have been generalized in [KZ10, KZ12]. In particular, in [KZ10] it has been showed that the f -Property implies an f -estimate for any f , and in [KZ12] that an f -estimate with $\frac{f}{\log} \rightarrow \infty$ at ∞ implies that the type along complex

analytic variety has a lower bound with the rate G with

$$G(\delta) = \left(\left(\frac{f}{\log} \right)^* (\delta^{-1}) \right)^{-1}, \quad (1.2)$$

where the superscript $*$ denotes the inverse function. Combining the above results, one obtains

Theorem 1.1 (Catlin [Cat83, Cat87]; Khanh-Zampieri [KZ10, KZ12]). *Let Ω be a pseudoconvex domain in $\mathbb{C}^n$ with C^∞ -smooth boundary and z_o be a point in the boundary $b\Omega$. Assume that the f -Property holds at z_o for $\frac{f}{\log} \nearrow \infty$ as $t \rightarrow \infty$ and that $b\Omega$ has type $\leq F$ along an one-dimensional complex analytic variety at z_o . Then, $F \gtrsim G$.*

The purpose of this note is to give an immediate proof of Theorem 1.1 which has also the advantage of requiring a minimal smoothness of $b\Omega$ and the expenses of a slightly stronger assumption about f . In fact, we prove the following

Theorem 1.2. *Let Ω be a pseudoconvex domain of $\mathbb{C}^n$ with C^2 -smooth boundary $b\Omega$ and z_o be a point in the boundary $b\Omega$. Assume that Ω has the f -Property at z_o with f satisfying $\int_t^\infty \frac{da}{af(a)} < \infty$ for some $t > 1$, denote by $(g(t))^{-1}$ this finite integral, and set $G(\delta) = (g^*(\delta^{-1}))^{-1}$. Then, if $b\Omega$ has type $\leq F$ along one-dimensional complex analytic variety Z , then $F(\delta) \gtrsim G(\alpha\delta)$ for a suitable α and for any δ small.*

First, we remark that the C^∞ -smoothness of the boundary in the results of Catlin and Khanh-Zampieri (Theorem 1.1) is required since they apply the regularity of the $\bar{\partial}$ -Neumann problem. In Theorem 1.2, the condition of smoothness is reduced because of the use of a plurisubharmonic peak function. However, in the construction of the family of the plurisubharmonic peak functions, we need a slightly stronger hypothesis in f , for example, if $f(t) = \log t \cdot \log^\epsilon(\log t)$ with $0 < \epsilon \leq 1$, then f satisfies the hypothesis in Theorem 1.1 but does not in Theorem 1.2. Finally, we notice the equivalence of the statements in the two theorems in case $f(t) = \log^\beta t$ for $\beta > 1$, or $f(t) = t^\epsilon$ for any $0 < \epsilon \leq \frac{1}{2}$.

2. PROOF OF THEOREM 1.2

The proof of Theorem 1.2 follows immediately from Theorem 2.1 and Theorem 2.2 below. In [Kha13], the author proves that there exists a family of plurisubharmonic functions with good estimates

Theorem 2.1. *Under the assumptions of Theorem 1.2, for a fixed constant $0 < \eta \leq 1$, there are a neighborhood V of z_o and positive constants c_1, c_2, c_3 such that the following holds. For any $w \in V \cap b\Omega$ there is a plurisubharmonic function ψ_w on $V \cap \Omega$ verifying*

$$(1) \quad |\psi_w(z) - \psi_w(z')| \leq c_1 |z - z'|^\eta,$$

- (2) $\psi_w(z) \leq -G^\eta(c_2|z - w|)$,
- (3) $\psi_{\pi(z)}(z) \geq -c_3\delta_{b\Omega}(z)^\eta$,

for any z and z' in $V \cap \bar{\Omega}$ (where $\delta_{b\Omega}(z)$ and $\pi(z)$ denote the distance and projection of z to the boundary, respectively).

Using the conclusion of Theorem 2.1 just for $w \equiv z_0$, we get

Theorem 2.2. *Let Ω be a C^2 -smoothly pseudoconvex domain in $\mathbb{C}^n$ and z_o be a boundary point. Assume that there is a neighborhood V of z_o and a plurisubharmonic function ψ on $V \cap \Omega$ such that*

$$-c_1|z - z_o|^\eta \leq \psi(z) \leq -G^\eta(c_2|z - z_o|), \quad z \in V \cap \Omega, \quad (2.1)$$

for suitable $c_1, c_2 > 0$ and $\eta \in (0, 1]$. In this situation, if $b\Omega$ has type $\leq F$ along a one-dimensional complex analytic variety Z , then $F(\delta) \gtrsim G(\alpha\delta)$.

In the case $G(t) = t^m$, the result has been obtained by Fornaess and Sibony in [FS89]. *Proof of Theorem 2.2.* Let Ω be a domain in $\mathbb{C}^n$ and assume that there is a function F and an 1-dimensional complex analytic variety Z passing through z_o such that (1.1) is satisfied for $z \in Z$. Then, in any neighborhood U of z_o there are constants $c_3, c_4 > 0$ and a family $\{Z_\delta\}$ of 1-dimensional complex manifolds $Z_\delta \subset U$ defined by $h_\delta : \bar{\Delta} \rightarrow U$ with $h_\delta(0) = z_o$ such that

$$\delta = \sup_{t \in \bar{\Delta}} |h_\delta(t) - z_o| \geq |h'_\delta(0)| \geq c_3\delta \quad (2.2)$$

and

$$\sup_{t \in \bar{\Delta}} |\delta_{b\Omega}(h_\delta(t))| < c_4F(\delta), \quad (2.3)$$

where Δ denotes the unit disc in $\mathbb{C}$.

Let ν be the outward normal to $b\Omega$ at z_o . From (2.3), we have $h_\delta(t) - c_4F(\delta)\nu \in \Omega \cap U$ for any $t \in \bar{\Delta}$. Applying the submean value inequality to the subharmonic function $\psi(h_\delta(t) - c_4F(\delta)\nu)$ on $\bar{\Delta}$ we get

$$\psi(z_o - c_4F(\delta)\nu) \leq \frac{1}{2\pi} \int_0^{2\pi} \psi(h_\delta(e^{i\theta}) - c_4F(\delta)\nu) d\theta. \quad (2.4)$$

Now, we use the first inequality in (2.1) for the left hand side term of (2.4)

$$-\psi(z_o - c_4F(\delta)\nu) \leq c_1c_4^\eta F^\eta(\delta)^\eta.$$

For the right hand side term of (2.4), we use the second inequality of (2.1)

$$-\frac{1}{2\pi} \int_0^{2\pi} \psi(h_\delta(e^{i\theta}) - c_4F(\delta)\nu) d\theta \geq \frac{1}{2\pi} \int_0^{2\pi} G^\eta(c_2|h_\delta(e^{i\theta}) - c_4F(\delta)\nu - z_o|) d\theta. \quad (2.5)$$

Using (2.2) and the Jensen inequality for the increasing, convex function G^η , we get

$$\begin{aligned} G^\eta(c_2 c_3 \delta) &\leq G^\eta(c_2 |h'_\delta(0)|) \leq G^\eta\left(\frac{1}{2\pi} \int_0^{2\pi} c_2 |h_\delta(e^{i\theta}) - CF(\delta)\nu - z_o| d\theta\right) \\ &\leq \frac{1}{2\pi} \int_0^{2\pi} G^\eta(c_2 |h_\delta(e^{i\theta}) - CF(\delta)\nu - z_o|) d\theta. \end{aligned} \quad (2.6)$$

Combining (2.4), (2.5) and (2.6), we obtain

$$F(\delta) \geq \beta G(\alpha \delta).$$

with $\beta = (c_1^{1/\eta} c_4)^{-1}$ and $\alpha = c_2 c_3$. The proof of Theorem (1.2) is complete. □

REFERENCES

- [Cat83] David Catlin. Necessary conditions for subellipticity of the $\bar{\partial}$ -Neumann problem. *Ann. of Math. (2)*, 117(1):147–171, 1983.
- [Cat87] D. Catlin. Subelliptic estimates for the $\bar{\partial}$ -Neumann problem on pseudoconvex domains. *Ann. of Math. (2)*, 126(1):131–191, 1987.
- [D'A82] John P. D'Angelo. Real hypersurfaces, orders of contact, and applications. *Ann. of Math. (2)*, 115(3):615–637, 1982.
- [FS89] J E. Fornaess and N. Sibony. Construction of P.S.H. functions on weakly pseudoconvex domains. *Duke Math. J.* 58 (3): 633–655, 1989.
- [Kha13] T. V Khanh. Boundary behavior of the Kobayashi metric near a point of infinite type. 2013.
- [KZ10] T. V Khanh and G. Zampieri. Regularity of the $\bar{\partial}$ -Neumann problem at point of infinite type. *J. Funct. Anal.*, 259(11):2760–2775, 2010.
- [KZ12] T. V Khanh and G. Zampieri. Necessary geometric and analytic conditions for general estimates in the $\bar{\partial}$ -Neumann problem. *Invent. Math.*, 188(3):729–750, 2012.